

## Stochastic Processes Sheet 1

Hand in Friday, 28th April 2017 before the lecture

### Exercise 1

[5 Pkt]

Find the open, closed and compact subsets of the metric space  $(\mathbb{Z}^d, \rho)$ , where  $\mathbb{Z}$  is the set of integer numbers and  $\rho$  is the euclidean metric in  $\mathbb{R}^d$ . Define the corresponding Borel- $\sigma$ -algebra  $\mathcal{B}(\mathbb{Z}^d)$ .

### Exercise 2

[5 Pkt]

Let  $\{(X_n, \mathcal{A}_n, \mu_n)\}_{n \in \mathbb{N}}$  be a family of measure spaces, where the sets  $X_n$  are pairwise disjoint. We define the measure space  $(X, \mathcal{A}, \mu)$ , where  $X = \bigcup_n X_n$ ,

$$\mathcal{A} = \{B : B \cap X_n \in \mathcal{A}_n \text{ for all } n\}$$

and  $\mu(B) = \sum_n \mu_n(B \cap X_n)$ . Show that

1.  $\mathcal{A}$  is a  $\sigma$ -algebra;
2.  $\mu$  is a measure;
3.  $\mu$  is  $\sigma$ -finite if and only if all  $\mu_n$  are  $\sigma$ -finite.

### Exercise 3

[5 Pkt]

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $A_1 \triangle A_2 = (A_1 \setminus A_2) \cup (A_2 \setminus A_1)$  for  $A_1, A_2 \in \mathcal{A}$ . Show that:

1. If  $A_1, A_2 \in \mathcal{A}$  and  $\mu(A_1 \triangle A_2) = 0$ , then  $\mu(A_1) = \mu(A_2)$ .
2. If the measure space is complete, from  $A_1 \in \mathcal{A}$  and  $\mu(A_1 \triangle A_2) = 0$ , it follows that  $A_2 \in \mathcal{A}$ .

*Remark:* A measure space  $(X, \mathcal{A}, \mu)$  is complete if the  $\sigma$ -algebra  $\mathcal{A}$  contains all the subsets of  $\mu$ -null sets, i.e., if  $B \in \mathcal{A}$ ,  $\mu(B) = 0$  and  $A \subset B$  then  $A \in \mathcal{A}$ .