

Sheet 5, “Markov Processes”

Due on November 20, 2018

Exercise 1

[5 Pt]

Define the family of operators $P_t, t \in \mathbb{R}_+$, on $C_0(\mathbb{R})$ by $P_t f(x) = f(x + t)$. Show that $P_t, t \in \mathbb{R}_+$, is a strongly continuous contraction semigroup and that its generator $(G, \mathcal{D}(G))$ is given by

$$Gf = f' \quad \text{and} \quad \mathcal{D}(G) = \{f \in C_0(\mathbb{R}) : f' \in C_0(\mathbb{R})\}.$$

Exercise 2

[5 Pt]

Let X be a one-dimensional Brownian motion. Define the family, $P_t, t \in \mathbb{R}_+$, of operators by $P_t f(x) := \mathbb{E}^x[f(X_t)]$ for $f \in C_0(\mathbb{R})$. Show that its generator $(G, \mathcal{D}(G))$ is given by

$$Gf = \frac{1}{2}f'' \quad \text{and} \quad \mathcal{D}(G) = \{f \in C_0(\mathbb{R}) : f', f'' \in C_0(\mathbb{R})\}.$$

Exercise 3

[5 Pt]

Let G be the linear operator on $C_0(\mathbb{R})$ defined by

$$Gf = f''' \quad \text{and} \quad \mathcal{D}(G) = \{f \in C_0(\mathbb{R}) : f', f'', f''' \in C_0(\mathbb{R})\}.$$

Show that G is *not* the generator of a strongly continuous contraction semi-group.

Hint: You may show that Theorem 4.19 (ii) is contradicted if you suppose that G would be the generator of a strongly continuous contraction semi-group.

Exercise 4

[5 Pt]

Let G be a closed operator on a Banach space B_0 , and let $\rho(G)$ be its resolvent set. Let $\lambda, \mu \in \rho(G)$. Show that the corresponding resolvents $R_\lambda = (\lambda - G)^{-1}$ and $R_\mu = (\mu - G)^{-1}$ satisfy the resolvent identity.