

Introduction to Stochastic Analysis, Problem sheet 8

Please hand in your solutions with your names and the name of your tutor
 before Tuesday 15.12., 12 am.

1. (Time-dependent Itô formula). Suppose that $X : [0, \infty) \rightarrow \mathbb{R}$ is a continuous function with continuous quadratic variation $[X]$ w.r.t. a sequence (π_n) of partitions s.t. $\text{mesh}(\pi_n) \rightarrow 0$. Show that for any function $F \in C^2(\mathbb{R}^2)$ and for any $t \in \mathbb{R}_+$, the Itô integral

$$\int_0^t \frac{\partial F}{\partial x}(s, X_s) dX_s = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} \frac{\partial F}{\partial x}(s, X_s) (X_{s' \wedge t} - X_{s \wedge t})$$

exists, and the time-dependent Itô formula

$$F(t, X_t) - F(0, X_0) = \int_0^t \frac{\partial F}{\partial s}(s, X_s) ds + \int_0^t \frac{\partial F}{\partial x}(s, X_s) dX_s + \frac{1}{2} \int_0^t \frac{\partial^2 F}{\partial x^2}(X_s) d[X]_s \quad (1)$$

holds.

Hint: You may assume without proof that by Taylor's formula, there exists a function $o : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $o(r)/r \rightarrow 0$ as $r \rightarrow 0$ such that for any $s, s' \in [0, t]$,

$$\begin{aligned} F(s', X_{s'}) - F(s, X_s) &= \frac{\partial F}{\partial s}(s, X_s) \delta s + \frac{\partial F}{\partial x}(s, X_s) \delta X_s + \frac{1}{2} \frac{\partial^2 F}{\partial s^2}(s, X_s) (\delta s)^2 \\ &\quad + \frac{\partial^2 F}{\partial s \partial x}(s, X_s) \delta s \delta X_s + \frac{1}{2} \frac{\partial^2 F}{\partial x^2}(s, X_s) (\delta X_s)^2 + o((\delta s)^2 + (\delta X_s)^2). \end{aligned}$$

2. (Geometric Brownian motion). Let (B_s) be an (\mathcal{F}_s) -Brownian motion on (Ω, \mathcal{A}, P) s.t. $B_0 = 0$. A geometric Brownian motion (X_s) with parameters $\mu, \alpha \in \mathbb{R}$ is a solution of the stochastic differential equation (SDE)

$$dX_t = \mu X_t dt + \alpha X_t dB_t,$$

i.e., (X_s) is an almost surely continuous and (\mathcal{F}_s) adapted process such that P -almost surely,

$$X_t - X_0 = \mu \int_0^t X_s ds + \alpha \int_0^t X_s dB_s \quad \text{for any } t \geq 0.$$

a) Find a solution of the SDE with initial value $X_0 = x_0$ using the ansatz

$$X_t = x_0 \cdot \exp(aB_t + bt).$$

Here you may assume the time-dependent Itô formula (1).

b) What can you say about the asymptotic behavior of the process as $t \rightarrow \infty$?

c) Compute $E[X_t]$ and $\text{Cov}[X_s, X_t]$ for $s, t \geq 0$.

3. (Stochastic integrals w.r.t. Itô processes). Let

$$I_s := \int_0^s H_r dB_r, \quad 0 \leq s \leq t,$$

with an (\mathcal{F}_s) -Brownian motion B on (Ω, \mathcal{A}, P) , and an (\mathcal{F}_s) -adapted process $H \in L^2(P \otimes \lambda)$. Suppose that (π_n) is a sequence of partitions of $[0, t]$ such that $\text{mesh}(\pi_n) \rightarrow 0$.

Prove that if G is another (\mathcal{F}_s) -adapted bounded continuous process, then the Riemann sums $\sum_{s \in \pi_n} G_s \cdot (I_{s'} - I_s)$ converge in $L^2(P)$, and

$$\int_0^t G_s dI_s = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} G_s \cdot (I_{s'} - I_s) = \int_0^t G_s H_s dB_s .$$

Hint: Express the Riemann sums as a stochastic integral $\int_0^t \dots dB_s$ w.r.t. Brownian motion.

4. (A local martingale that is not a martingale). Let $(B_t)_{t \geq 0}$ be a Brownian motion in \mathbb{R}^3 with initial value $B_0 = x$, $x \neq 0$. Show that:

- a) $X_t = 1/\|B_t\|$ is a local martingale up to $T = \infty$ w.r.t. the filtration generated by (B_t) .
- b) $\{X_s : 0 \leq s \leq t\}$ is uniformly integrable for all $t \geq 0$.
- c) X_t is *not* a martingale.

Hint: You may assume without proof the multi-dimensional Itô formula for Brownian motion: If U is an open subset of \mathbb{R}^d then for $F \in C^2(U)$,

$$F(B_t) - F(B_0) = \int_0^t \nabla F(B_s) \cdot dB_s + \frac{1}{2} \int_0^t \Delta F(B_s) ds \quad \forall t < T_{UC} .$$