

„Markov Processes”, Problem Sheet 1

Please hand in your solutions before 12 am on Monday, October 14,
into the marked post boxes opposite to the maths library.

1. (Conditional Expectations).

- a) Let X, Y on (Ω, \mathcal{A}, P) be independent random variables that are Bernoulli distributed with parameter p . We set $Z = 1_{\{X+Y=0\}}$. Compute $E[X|Z]$ and $E[Y|Z]$. Are these random variables still independent?
- b) Let X, Y, Z be random variables with values in a measurable space (S, \mathcal{B}) such that the couples (X, Z) and (Y, Z) have the same law. Show that, for any $f \in \mathcal{F}_+(S)$,

$$E[f(X)|Z] = E[f(Y)|Z] \quad \text{a.s.}$$

- c) Let T_1, \dots, T_n be i.i.d. real integrable random variables. Set $T = T_1 + \dots + T_n$.
- (i) Show that $E[T_1|T] = T/n$.
- (ii) Compute $E[T|T_1]$.

2. (Conditional laws of Gaussian vectors).

- a) Let X be a random variable with values in \mathbb{R}^m of the form $X = \phi(Y) + Z$, where Y and Z are independent. Show that the conditional law of X given $Y = y$ coincides with the law of $\phi(y) + Z$.
- b) Let X, Y be random variables taking values in \mathbb{R}^k and \mathbb{R}^p , respectively. We assume that their joint law in $(\mathbb{R}^{k+p}, \mathcal{B}(\mathbb{R}^{k+p}))$ is Gaussian with mean and covariance matrix given, respectively, by

$$\begin{pmatrix} m_X \\ m_Y \end{pmatrix}, \quad \begin{pmatrix} R_X & R_{XY} \\ R_{YX} & R_Y \end{pmatrix}.$$

Here m_X, m_Y , and R_X, R_Y are the means and covariance matrices of X and Y , respectively, and $R_{X,Y} = E[(X - E[X])(Y - E[Y])^T] = R_{Y,X}^T$ is the $k \times p$ matrix of the covariances between components of X and Y . We assume that R_Y is strictly positive definite.

- (i) Find a $k \times p$ matrix A such that the random variables $X - AY$ and Y are independent.
- (ii) Show that the conditional law of X given Y is Gaussian with mean $E[X|Y] = m_X + R_{XY}R_Y^{-1}(Y - m_Y)$ and covariance matrix $R_X - R_{XY}R_Y^{-1}R_{YX}$.

- c) Let X be a signal with normal law $\mathcal{N}(0, 1)$. We assume that we cannot observe the value of X ; instead we only observe the value of $Y = X + W$, where W is independent of X with law $\mathcal{N}(0, \sigma^2)$.
- (i) Given an observation $Y = y$, give an estimate of the value X of the signal.
 - (ii) Let us assume that $\sigma^2 = 0.1$ and that the value of the observation is $Y = 0.55$. What is the probability for the signal X to be in the interval $[\frac{1}{4}, \frac{3}{4}]$?

3. (Martingales with independent increments).

A process $(M_n)_{n \in \mathbb{Z}_+}$ is said to be with independent increments if, for every n , the r.v. $M_{n+1} - M_n$ is independent of the σ -algebra $\mathcal{F}_n = \sigma(M_0, \dots, M_n)$.

- a) Let $(M_n)_{n \in \mathbb{Z}_+}$ be a square integrable martingale with independent increments. We set $\sigma_0^2 = \text{Var}(M_0)$ and, for $k \geq 1$, $\sigma_k^2 = \text{Var}(M_k - M_{k-1})$.
 - (i) Show that $\text{Var}(M_n) = \sum_{k=1}^n \sigma_k^2$.
 - (ii) Determine the conditional variance process $\langle M \rangle_n$ of (M_n) .
- b) Let (M_n) be a Gaussian martingale (we recall that a process $(M_n)_{n \geq 0}$ is Gaussian if, for every n , the vector (M_0, \dots, M_n) is Gaussian).
 - (i) Show that $(M_n)_{n \geq 0}$ has independent increments.
 - (ii) Show that, for every fixed $\theta \in \mathbb{R}$, the process $Z_n^\theta = \exp(\theta M_n - \frac{1}{2}\theta^2 \langle M \rangle_n)$ is a martingale. Does it converge almost surely?