

„Markov Processes”, Problem Sheet 7

On this sheet, Exercise 2 counts for two exercises.

Please hand in your solutions before 12 noon on Monday, November 25.

1. (Equivalent characterizations of ergodicity for Markov processes). We consider a canonical right-continuous Markov process $((X_t)_{t \geq 0}, (P_x)_{x \in S})$ with Polish state space (S, \mathcal{B}) , transition semigroup $(p_t)_{t \geq 0}$, and invariant probability measure μ .

a) Prove that the following nine conditions are all equivalent:

- (i) P_μ is ergodic.
- (ii) $\frac{1}{t} \int_0^t F \circ \Theta_s ds \rightarrow E_\mu[F]$ P_μ -a.s. and in $L^2(P_\mu)$, for any $F \in \mathcal{L}^2(P_\mu)$.
- (iii) $\frac{1}{t} \int_0^t f(X_s) ds \rightarrow \int f d\mu$ P_μ -a.s. and in $L^2(P_\mu)$, for any $f \in \mathcal{L}^2(\mu)$.
- (iv) $\frac{1}{t} \int_0^t \text{Cov}_{P_\mu} [f(X_0), g(X_s)] ds \rightarrow 0$ as $t \uparrow \infty$ for any $f, g \in \mathcal{L}^2(\mu)$.
- (v) $\frac{1}{t} \int_0^t P_\mu[X_0 \in B, X_s \in C] ds \rightarrow \mu(B)\mu(C)$ for any $B, C \in \mathcal{B}$.
- (vi) $\frac{1}{t} \int_0^t p_s(x, B) ds \rightarrow \mu(B)$ μ -a.e. for any $B \in \mathcal{B}$.
- (vii) $P_x[\exists t \in \mathbb{Q}_+ : X_t \in B] > 0$ μ -a.e. for any $B \in \mathcal{B}$ such that $\mu(B) > 0$.
- (viii) Every set $B \in \mathcal{B}$ such that $p_t 1_B = 1_B$ μ -a.e. for any $t \geq 0$ satisfies $\mu(B) \in \{0, 1\}$.
- (ix) Every function $h \in \mathcal{L}^2(\mu)$ such that $p_t h = h$ μ -a.e. $\forall t \geq 0$ is μ -almost surely constant.

b) A function $h : S \rightarrow \mathbb{R}$ is called *harmonic* w.r.t. (p_t) iff $p_t h = h$ for any $t \geq 0$. Show that the following statements are all equivalent:

(i) For every $x \in S$ and $B \in \mathcal{B}$ with $\mu(B) > 0$,

$$P_x [\forall s > 0 \exists t \geq s : X_t \in B] = 1.$$

(ii) The constants are the only bounded harmonic functions for (p_t) .

(iii) For every bounded measurable function $F : \Omega \rightarrow \mathbb{R}$ and every $x \in S$,

$$\frac{1}{t} \int_0^t F \circ \Theta_s ds \rightarrow E_\mu[F] \quad P_x\text{-almost surely.}$$

2. (Brownian motion on \mathbb{R}/\mathbb{Z}). A Brownian motion (X_t) on the circle \mathbb{R}/\mathbb{Z} can be obtained by considering a Brownian motion (B_t) on \mathbb{R} modulo the integers, i.e.,

$$X_t = B_t - [B_t] \in [0, 1) \cong \mathbb{R}/\mathbb{Z}.$$

Prove the following statements:

- a) Brownian motion on \mathbb{R}/\mathbb{Z} is a Markov process with transition density w.r.t. the uniform distribution given by

$$p_t(x, y) = \frac{1}{\sqrt{2\pi t}} \sum_{n \in \mathbb{Z}} e^{-\frac{|x-y-n|^2}{2t}} \quad \text{for any } t > 0 \text{ and } x, y \in [0, 1).$$

- b) For any initial condition, (X_t) solves the martingale problem for the operator $\mathcal{L}f = f''/2$ defined on $C^\infty(\mathbb{R}/\mathbb{Z})$. (Note that there is a one-to-one correspondence of smooth functions on \mathbb{R}/\mathbb{Z} and periodic smooth functions on \mathbb{R} with period 1).
- c) The uniform distribution μ is an invariant probability measure for (p_t) , and the process with initial distribution μ is stationary and ergodic.
- d) The generator $-\mathcal{L}$ has smooth real-valued eigenfunctions e_n , $n \in \mathbb{Z}$, with corresponding eigenvalues $\lambda_n = 2\pi^2 n^2$. Moreover, $p_t e_n = \exp(-\lambda_n t) e_n$ for any $t \geq 0$.
- e) For any $f \in \mathcal{L}^2(\mu)$,

$$\left\| p_t f - \int f d\mu \right\|_{L^2(\mu)} \leq e^{-2\pi^2 t} \sqrt{\text{Var}_\mu(f)}.$$

- f) Conclude that for the process with initial distribution μ ,

$$E \left[\left(\frac{1}{t} \int_0^t f(X_s) ds - \int f d\mu \right)^2 \right] \leq \frac{1}{\pi^2 t} \text{Var}_\mu(f) \quad \text{for any } t \geq 0 \text{ and } f \in \mathcal{L}^2(\mu).$$

3. (Metropolis-Hastings method). Let $\mu(dx) = \mu(x) dx$ be a probability measure on \mathbb{R}^d with strictly positive density, and let $q(x, dy) = q(x, y) dy$ be a probability kernel on \mathbb{R}^d with strictly positive density. The Metropolis-Hastings acceptance probability is given by

$$\alpha(x, y) = \min \left(1, \frac{\mu(y)q(y, x)}{\mu(x)q(x, y)} \right), \quad x, y \in \mathbb{R}^d.$$

Metropolis-Hastings algorithm

- 1.) Set $n := 0$ and choose some arbitrary point $X_0 \in \mathbb{R}^d$.
- 2.) Sample $Y_{n+1} \sim q(X_n, \cdot)$ and $U_{n+1} \sim \text{Unif}(0, 1)$ independently.
- 3.) If $U_{n+1} < \alpha(X_n, Y_{n+1})$ then set $X_{n+1} := Y_{n+1}$, else set $X_{n+1} := X_n$.
- 4.) Set $n := n + 1$ and go to Step 2.

Show that for every bounded measurable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\frac{1}{n} \sum_{i=0}^{n-1} f(X_i) \rightarrow \int f d\mu \quad \text{almost surely.}$$

The student council of mathematics will organize the Maths Party on 28/11 in N8schicht. The presale will be held on Tue 26/11, Wed 27/11 and Thu 28/11 in the mensa Poppelsdorf. Further information can be found on fsmath.uni-bonn.de.