

## Sheet 2, “Stochastic Analysis”

To hand in until April 17, 15:00

---

Check the exercise classes on ecampus for the hand-in. You should submit your solutions in groups of two, preferably (but not necessarily) with another student from the same tutorial. Please upload only compressed PDFs of decent quality.

---

### Problem 1 (Scale function: recurrence and exit times, 7 Pt)

Suppose that  $(X_t)_{0 \leq t \leq \xi}$  is a strong solution of the SDE

$$\begin{aligned}dX_t &= b(X_t)dt + \sigma(X_t)dB_t, \quad t \in [0, \xi), \\ X_0 &= x_0,\end{aligned}$$

for a one-dimensional Brownian motion  $B_t$ , an initial value  $x_0 \in (0, \infty)$  and continuous  $b, \sigma \in (0, \infty)$ . We fix the left boundary and assume that the solution exists up to the explosion time

$$\xi := \sup_{\epsilon, r > 0} T_{\epsilon, r}, \quad T_{\epsilon, r} = \inf\{t \geq 0 \mid X_t \notin (\epsilon, r)\}.$$

The scale-function  $s : (0, \infty) \rightarrow \mathbb{R}$  of this process is given by

$$s(x) := \int_{x_0}^x \exp \left[ - \int_{x_0}^z \frac{2b(y)}{\sigma(y)^2} dy \right] dz.$$

One can easily show that for  $\epsilon < x_0 < r$ :

$$\mathbb{P}[T_\epsilon < T_r] = \frac{s(r) - s(x_0)}{s(r) - s(\epsilon)}.$$

In the following, assume that  $s(x)$  converges as  $x \rightarrow 0$  and  $x \rightarrow \infty$ .

1) Conclude that

1. If  $s(0) = -\infty$  and  $s(\infty) = \infty$ , then  $X_t$  is recurrent. For any  $x_0, y \in (0, \infty)$ :

$$\mathbb{P}[X_t = y \text{ for some } t \in [0, \xi)] = 1.$$

2. If  $s(0) > -\infty$  and  $s(\infty) < \infty$ , then

$$\mathbb{P} \left[ \lim_{t \rightarrow \xi} X_t = 0 \right] = \frac{s(\infty) - s(x_0)}{s(\infty) - s(0)}.$$

3. If  $s(0) > -\infty$  and  $s(\infty) = \infty$ , then  $\lim_{t \rightarrow \xi} X_t = 0$  a.s. .

4. If  $s(0) = -\infty$  and  $s(\infty) < \infty$ , then  $\lim_{t \rightarrow \xi} X_t = \infty$  a.s. .

2) Let  $\alpha \in \mathbb{R}$  and  $\sigma > 0$ . Let  $(S_t)_{0 \leq t < \xi}$  be a Geometric Brownian motion, a strong solution of the SDE

$$dS_t = \alpha S_t dt + \sigma S_t dB_t, \quad S_0 = x_0 > 0.$$

Analyze the asymptotic behavior of  $S_t$ .

3) Let  $\beta \in \mathbb{R}$  and  $\sigma > 0$ . Let  $(F_t)_{0 \leq t < \xi}$  be a strong solution of Feller's branching diffusion:

$$dF_t = \beta F_t dt + \sigma \sqrt{F_t} dB_t, \quad F_0 = x_0 > 0.$$

Analyze the asymptotic behavior of  $F_t$ .

*Remark:* The process  $F_t$  is the diffusive limit of a discrete Galton-Watson branching process, describing a population in which the number  $N$  of offsprings of each individual has the following properties:  $\mathbb{E}[N] = 1 + \beta$ ,  $\text{Var}[N] = \sigma^2$ . Hence,  $\mathbb{P}[F(\xi) = 0]$  is the probability that this population dies out.

### **Problem 2 (Local time and mode of continuity of Brownian motion, 3 Pt)**

Let  $(B_t)_{t \geq 0}$  be a one-dimensional standard Brownian motion and let  $\Gamma_t$  be the occupation measure as defined in the lecture. Its density  $l_t^x$  is jointly continuous in  $t$  and  $x$  almost surely. Show that for almost all  $\omega$ , the path  $t \mapsto B_t(\omega)$  is not locally Lipschitz continuous.

*Hint:* Consider  $\Gamma_{t+h}(S_h) - \Gamma_t(S_h)$  for suitable sets  $S_h$ .